

PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Keep away from humans, domestic animals and pets. May be harmful or fatal if swallowed or absorbed through the skin because this material may reduce the clotting ability of blood and cause bleeding.

Do not get in eyes, on skin or on clothing. Wash arms, hands and face with soap and water after applying and before eating and smoking. **If swallowed, call a physician at once.**

NOTE TO PHYSICIAN: If ingested, administer vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds.

STORAGE AND DISPOSAL:

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. STORE IN ORIGINAL CONTAINER IN LOCKED STORAGE AREA.

STORAGE: Store in original container in a cool, dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Securely wrap excess, unused or spoiled bait in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse pail. Rinse thoroughly before discarding in trash.

KEEP AWAY FROM FOODSTUFFS

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA REG. NO. 7173-188

EPA EST. NO 7173-WI-1



PARAFFINIZED PELLETS

FOR INDOOR AND OUTDOOR RODENT CONTROL

KILLS NORWAY RATS, ROOF RATS, HOUSE MICE AND WARFARIN RESISTANT NORWAY RATS

NORWAY RATS AND HOUSE MICE MAY CONSUME A LETHAL DOSE IN ONE FEEDING WITH FIRST DEAD RODENTS APPEARING FOUR OR FIVE DAYS AFTER TREATMENT BEGINS.

ACTIVE INGREDIENT:

*3-[3-(4'-bromo[1,1'-biphenyl]-4-yl)-3-hydroxy-1-phenylpropyl]-4-hydroxy-2H-1-benzopyran-2-one.....	0.005%
Inert ingredients.....	99.995%
TOTAL.....	100.000%

*Bromadiolone-Supercaid Patented in the U.S.A.

KEEP OUT OF REACH OF CHILDREN CAUTION

MAY BE HARMFUL OR FATAL IF SWALLOWED. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.



Manufactured By
LiphaTech Inc.
3101 W. Custer Ave.
Milwaukee, WI 53209

Net Weight: _____

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets or other nontarget animals to rodenticides. To help prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hooved livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container, unused, spoiled and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of Norway rats, roof rats and house mice.

URBAN AREAS: This product may be used in and around the periphery of homes, industrial, commercial and public buildings. This product may also be used in and around transport vehicles (ships, trains, aircraft) and related port or terminal buildings. This product may be used in alleys.

NONURBAN AREAS: This product may be used inside of homes and agricultural buildings.

Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. Do not broadcast bait.

SELECTION OF TREATMENT AREAS:

Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

RATS: Place at least 3 to 10 packs per placement. Maintain an uninterrupted supply of fresh bait for 10 days or until signs of rat activity cease.

MICE: Use one pack per bait placement. Space placements at 8- to 12-foot intervals. Maintain an uninterrupted supply of fresh bait for 15 days or until fresh signs of mouse activity cease to appear.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.